

mood-book



Test UNIT-IV

Large sample test (n > 30)

1) Test of significance of single mean

\bar{x}, n

Table value = $Z_{\alpha/2}$ (or) Z_{α}

2) Test of sign. difference b/w two means

$\bar{x} \quad \bar{y}$
 $n_1 \quad n_2$
 $\sigma_1 \quad \sigma_2$

3) Test of sign. of single proportion

$\bar{x}, \sigma \rightarrow$ not given.

$p = \frac{x}{n}$

4) Test of sign. difference b/w two proportions.

$\bar{x}, \sigma \rightarrow$ not given.

$x_1 \quad x_2$
 $n_1 \quad n_2$

$P_1 = \frac{x_1}{n_1} \quad P_2 = \frac{x_2}{n_2}$

Small Sample Test (n < 30)

1) t-test

Test of significance of single mean.

$\bar{x}, n, v = (n-1)$

$S^2 = \frac{1}{n-1} \sum (x_i - \bar{x})^2$

Test Statistic.

$z = \frac{\bar{x} - \mu}{\sigma/\sqrt{n}}$ $H_0: \mu = \mu_0$
 $H_1: \mu \neq \mu_0$

$z = \frac{\bar{x} - \bar{y}}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$ $H_0: \mu_1 = \mu_2$
 $H_1: \mu_1 \neq \mu_2$

$z = \frac{p - P}{\sqrt{\frac{PQ}{n}}}$ $H_0: P = P_0$
 $H_1: P \neq P_0$

$Q = 1 - P$

$z = \frac{p_1 - p_2}{\sqrt{p_0(\frac{1}{n_1} + \frac{1}{n_2})}}$ $H_0: P_1 = P_2$
 $H_1: P_1 \neq P_2$

where $p =$ combined proportion.
 $p = \frac{n_1 p_1 + n_2 p_2}{n_1 + n_2} = \frac{x_1 + x_2}{n_1 + n_2}$

$Q = 1 - p$

case i) if 's' is given

$t = \frac{\bar{x} - \mu}{s/\sqrt{n-1}}$

case ii) if s is calculated (s) i.e. s is not given directly

$t = \frac{\bar{x} - \mu}{S/\sqrt{n}}$

for two tailed:
Table value = $t_{\alpha/2}$

for one-tailed:
Table value = t_{α}

2) Test of sign. difference b/w two means.

\bar{x}	\bar{y}
n_1	n_2
s_1^2	s_2^2

for t degrees of freedom:
 $v = n_1 + n_2 - 2$

Two tailed test:-
Table value = $t_{\alpha/2}$

one tailed test:-
Table value = t_{α}

3) paired t -test:-

x_i	Before	
y_i	After	

$$d_i = x_i - y_i \quad (\text{or}) \quad d_i = y_i - x_i$$

$$d.f (v) = (n - 1)$$

F-test

Case 1) If s_1, s_2 are given directly.

$$s^2 = \frac{n_1 s_1^2 + n_2 s_2^2}{n_1 + n_2 - 2}$$

$$t = \frac{\bar{x} - \bar{y}}{s \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

Case 2) If s_1, s_2 are not given directly.

$$s^2 = \frac{1}{n_1 + n_2 - 2} \left[\sum (x_i - \bar{x})^2 + \sum (y_j - \bar{y})^2 \right]$$

$$\bar{x} = \frac{1}{n_1} \sum x_i \quad \bar{y} = \frac{1}{n_2} \sum y_j$$

$$t = \frac{\bar{x} - \bar{y}}{s \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

$$t = \frac{\bar{d}}{s/\sqrt{n}}$$

$$\bar{d} = \frac{1}{n} \sum d_i$$

$$s^2 = \frac{1}{n-1} \sum (d_i - \bar{d})^2$$

F-test

Variances

$$H_0: \sigma_1^2 = \sigma_2^2$$

$$H_1: \sigma_1^2 \neq \sigma_2^2$$

Table Value

$$F_{\alpha}(V_1, V_2)$$

$$V_1 = (n_1 - 1)$$

$$V_2 = (n_2 - 1)$$

χ^2 -test

1) Goodness of Fit

$$H_0: O_i = E_i$$

$$H_1: O_i \neq E_i$$

Table Value

$$d.f. (v) = n - 1$$

$$\chi^2_{\alpha}$$

2) Independence of attributes

H_0 : there is no association b/w attributes.

H_1 : There is an association b/w attributes.

	C	D	
A	a	b	a+b
B	c	d	
	a+c	b+d	a+b+c+d

\Rightarrow

A	$\frac{(a+b)(a+c)}{N}$	$\frac{(b+d)(a+c)}{N}$	a+b
B	$\frac{(a+c)(b+d)}{N}$	$\frac{(b+d)(b+d)}{N}$	c+d

$N = a+b+c+d$

$$\chi^2 = \sum \frac{(O_i - E_i)^2}{E_i}$$

$$d.f. (v) = (\text{No. of rows} - 1) \times (\text{No. of columns} - 1)$$

$$\text{Table value} = \chi^2_{\alpha}$$

$$F = \frac{\text{Greater variance}}{\text{Smaller variance}}$$

$$F = \frac{S_1^2}{S_2^2}, \text{ if } S_1^2 > S_2^2$$

$$S_1^2 = \frac{n_1 S_1'^2}{n_1 - 1} = \frac{\sum (x_i - \bar{x})^2}{n_1 - 1}$$

$$S_2^2 = \frac{n_2 S_2'^2}{n_2 - 1} = \frac{\sum (y_j - \bar{y})^2}{n_2 - 1}$$

$$\chi^2 = \sum \frac{(O_i - E_i)^2}{E_i}$$

UNIT - V

Stochastic process (or) Random process :-

A set of Random variable values $\{x_t\}$ (or) $\{x_i\}$ depending on some real parameters (time t , temperature etc.) is known as stochastic process.

States :-

The values assumed by R.V.

State space :-

The set of all possible values of any individual no. of random process is called state space.

It is denoted by $\{x_t, t \in T\} = I$ (or) S .

Ex When a fair die is tossed, the no. of sixes is stochastic process

If the parameter set is discrete then the state space is discrete.

If the parameter set has infinite values, the state space is continuous.

* * Classification of stochastic process :-

1)

		time	
R.V		x	t
continuous	continuous	continuous stochastic process	continuous stochastic sequence
	discrete	discrete stochastic process	discrete stochastic sequence

Stationary stochastic process:

If the probability distribution do not depends on the time 't' then the random process is called stationary stochastic process.

Deterministic stochastic process:-

A Random process is called deterministic stochastic process if future values of any sample functions can be predicted from its past observations.

Non-Deterministic

A stochastic process is called non deterministic if future values of any sample functions can't be predicted from its past observations

Markov process:-

A Random process X_n is called markov process

if $P\{X(t_{n+1}) < X_{n+1} | X(t_n) < X_n, X(t_{n-1}) < X_{n-1}, \dots, X(t_0) < X_0\}$,

$$\Rightarrow P\{X(t_{n+1}) < X_{n+1} | X(t_n) < X_n\}$$

$$\Rightarrow P(X_{n+1} | X_n)$$

$X_{n+1}, X_n, X_{n-1}, X_{n-2}, \dots, X_0$

All states of micro processor

(or)

$$P\{X_n = k | X_{n-1} = j, X_{n-2} = i, \dots, X_0 = j_{n-1}\}$$

$$\Rightarrow P\{X_n = k | X_{n-1} = j\}$$

$$\Rightarrow P_{jk}^{(n)}$$

unit step transition probability :-

The probability $P_{jk}^{(1)}$ is called unit step transition probability ,

M-step transition probability :-

$$P \{ X_{n+m} = k \mid X_n = j \} = P_{jk}^{(m)}$$

Homogeneous Markov process :-

m-step If the transition probability P_{jk} is independent of 'n', then the Markov chain is called homogeneous Markov process :-

Non-Homogeneous

If the transition probability P_{jk} is dependent of 'n' then the step is called Markov non-homogeneous Markov process.

Probability distribution vector :-

A row or column matrix which consists of the probabilities of occurrences of Markov process then it is known as probability distribution vector.

If $P_1, P_2, P_3, \dots, P_n$ are probabilities.

then it is $[P_1, P_2, \dots, P_n]$

Transition probability matrix :-

The transition probabilities P_{jk} satisfies

i) $P_{jk} \geq 0$ i.e., non-negative elements.

ii) $\sum_{k=1}^n P_{jk} = 1 \quad \forall j$. i.e., each row sum = 1

$$P = [P_{jk}]_{m \times n} = \begin{bmatrix} P_{11} & P_{12} & \dots & P_{1n} \\ P_{21} & P_{22} & \dots & P_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ P_{m1} & P_{m2} & \dots & P_{mn} \end{bmatrix}$$

This is called transition probability matrix.

Stochastic matrix:-

A transition probability matrix is called stochastic matrix if it is square matrix.

Regular matrix:-

A tpm is called regular matrix if it satisfies

- i) stochastic matrix
- ii) diagonal element shouldn't equal to 1.
- iii) All elements of P^m , $m = 2, 3, \dots$ are positive.

Which of the following matrices are stochastic matrix

i) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ii) $\begin{bmatrix} 1/2 & 1/2 \\ 1/2 & 1/2 \end{bmatrix}$ iii) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$

iv) $\begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix}$ v) $\begin{bmatrix} 0 & 2 \\ 1/4 & 1/4 \end{bmatrix}$

i) stochastic

ii) stochastic

iii) Not stochastic. It is rectangular matrix.

iv) not stochastic. Negative element.

v) Row sum $\neq 1$

\therefore Non stochastic

2) which of the following are regular matrices.

$$i) A = \begin{bmatrix} 1/2 & 1/4 & 1/4 \\ 0 & 1 & 0 \\ 1/2 & 0 & 1/2 \end{bmatrix}$$

$$ii) B = \begin{bmatrix} 1/2 & 1/2 & 0 \\ 1/2 & 1/2 & 0 \\ 1/4 & 1/4 & 1/2 \end{bmatrix}$$

$$iii) C = \begin{bmatrix} 0 & 0 & 1 \\ 1/2 & 0 & 1/2 \\ 0 & 1 & 0 \end{bmatrix}$$

$$iv) D = \begin{bmatrix} 1/2 & 1/4 & 1 \\ 0 & 1/2 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

i) It is not regular, since, 1 lies on the diagonal.

$$ii) B^2 = B \cdot B = \begin{bmatrix} 1/2 & 1/2 & 0 \\ 1/2 & 1/2 & 0 \\ 1/4 & 1/4 & 1/2 \end{bmatrix} \begin{bmatrix} 1/2 & 1/2 & 0 \\ 1/2 & 1/2 & 0 \\ 1/4 & 1/4 & 1/2 \end{bmatrix} = \begin{bmatrix} 1/2 & 1/2 & 0 \\ 1/2 & 1/2 & 0 \\ 3/8 & 3/8 & 0 \end{bmatrix}$$

$$B^3 = B^2 \cdot B = \begin{bmatrix} 1/2 & 1/2 & 0 \\ 1/2 & 1/2 & 0 \\ 3/8 & 3/8 & 0 \end{bmatrix} \begin{bmatrix} 1/2 & 1/2 & 0 \\ 1/2 & 1/2 & 0 \\ 1/4 & 1/4 & 1/2 \end{bmatrix} = \begin{bmatrix} 1/2 & 1/2 & 0 \\ 1/2 & 1/2 & 0 \\ 7/16 & 7/16 & 1/8 \end{bmatrix}$$

The elements B_{13}, B_{23} are non-zeroes.

$$iii) C^3 = C^2 \cdot C = \begin{bmatrix} 1/2 & 0 & 1/2 \\ 1/4 & 1/2 & 1/4 \\ 0 & 1/2 & 1/2 \end{bmatrix}$$

$$C^4 = C^3 \cdot C = \begin{bmatrix} 0 & 1/2 & 1/2 \\ 1/4 & 1/4 & 1/2 \\ 1/4 & 1/2 & 1/4 \end{bmatrix}$$

$$C^5 = C^4 \cdot C = \begin{bmatrix} 1/4 & 1/2 & 1/4 \\ 1/8 & 1/2 & 3/8 \\ 1/4 & 1/4 & 1/2 \end{bmatrix}$$



The all elements of C^5 is non-zero.

\therefore It is regular stochastic matrix.

iv) Not a regular stochastic matrix.

3) Represent the following matrices as a transition matrices as a digraph.

i)
$$\begin{bmatrix} 0.75 & 0.75 & 0 \\ 0 & 1 & 0 \\ 0.3 & 0.5 & 0.2 \end{bmatrix}$$
 ii)
$$\begin{matrix} 1 & 2 & 3 \\ \begin{bmatrix} 3/4 & 1/4 & 0 \\ 1/4 & 1/2 & 1/4 \\ 0 & 3/4 & 1/4 \end{bmatrix} \end{matrix}$$



Steady state prob. Distribution :-

If $p^{(0)}$ is initial state probability distribution vector then after 2 step the distribution vector becomes $p^{(1)} = p^{(0)} \cdot P$. Here $P = TPM$.

After 2 steps :-

$$p^{(2)} = p^{(1)} \cdot P = p^{(0)} \cdot P^2$$

After n steps :-

$$p^{(n)} = p^{(n-1)} P = p^{(0)} \cdot P^n$$

If a homogeneous markov chain is regular, every sequence of state probability distribution approaches a unique fixed prob. distribution is called stationary distribution (or) steady state distribution of markov chain.

When, $n \rightarrow \infty$

$$\lim_{n \rightarrow \infty} p^{(n)} = \pi$$

$$\text{where } \pi = [\pi_1, \pi_2, \dots, \pi_n]$$

and the steady state distribution satisfies

$$\underline{\pi P = \pi} \text{ and } \pi_1 + \pi_2 + \dots + \pi_n = 1$$

Chapman - Kolmogorov's Theorem:-

If P is tpm of homogeneous Markov chain then n step tpm ($P^{(n)}$) is equal to P^n .

$$\left[P_{ij}^{(n)} \right] = \left[P_{ij} \right]^n$$

Classification of states:-

i) Irreducible:-

If $P_{ij}^{(n)} > 0$ for some 'n' and for every i and j , then every state can be reached from every other state.

ie, All states are communicate themselves.

Then the Markov chain is said to be irreducible.

The tpm of irreducible chain is called irreducible matrix, otherwise it is reducible.

ii) The state i of Markov chain is called recurrent state if $P_{ii}^{(n)} > 0$ for some 'n'.

iii) The period d_i of recurrent state i is defined as the greatest common divisor of all m there exists $P_{ii}^{(m)} > 0$,

$$d_i = \text{GCD} \{ m, P_{ii}^{(m)} > 0 \} = d_i$$

The state i is said to be periodic if $d_i > 1$ and aperiodic if $d_i = 1$.

v) The state 'i' is aperiodic if $p_{ii} \neq 0$

iv) The probability that the chain returns to state 'i' having started from 'i', for the first time at n^{th} step is denoted by $f_{ii}^{(n)}$, it is called first return time prob, $\{n, f_{ii}^{(n)}, n=1, 2, 3, \dots\}$ is called the distribution recurrence time state 'i',

$$F_{ii} = \sum_{n=1}^{\infty} f_{ii}^{(n)}$$

$$u_{ii} = \sum_{n=1}^{\infty} n \cdot f_{ii}^{(n)}$$

if $F_{ii} = 1$, then the state 'i' is called recurrent (or) persistent or return to the state 'i' it is also called certain.

If $F_{ii} < 1$, then the state 'i' is called transient or uncertain.

u_{ii} is called mean recurrence time of state 'i'.

If u_{ii} is finite then it is called non-null persistent.

If u_{ii} is infinite then it is called null persistent.

v) A non null persistent and aperiodic state is called ergodic.

vi) If a Markov chain is finite irreducible then all its states are non-null persistent.

$$\begin{aligned} \text{vii) } P(x_3=a, x_2=b, x_1=c, x_0=a) &= P(x_3=a/x_2=b) P(x_2=b/x_1=c) \\ &\quad P(x_1=c/x_0=a) \cdot P(x_0=a) \end{aligned}$$

1) A Markov

1) If the tpm of a Markov chain is $\begin{bmatrix} 0 & 1 \\ 1/2 & 1/2 \end{bmatrix}$ then find steady state distribution of chain.

Let $\pi = [\pi_1, \pi_2]$ be steady state distr. vector.

$$\pi P = \pi, \quad \pi_1 + \pi_2 = 1 \rightarrow (1)$$

$$[\pi_1 \ \pi_2] \begin{bmatrix} 0 & 1 \\ 1/2 & 1/2 \end{bmatrix} = [\pi_1 \ \pi_2]$$

$$\left[\frac{1}{2} \pi_2 \quad \pi_1 + \frac{1}{2} \pi_2 \right] = [\pi_1 \ \pi_2]$$

compare same position elements.

$$\frac{1}{2} \pi_2 = \pi_1, \quad \pi_1 + \frac{1}{2} \pi_2 = \pi_2$$

$$\pi_2 = 2\pi_1 \rightarrow (2) \quad 2\pi_1 + \pi_2 = 2\pi_2 \rightarrow (3)$$

from (2), (1).

$$\pi_1 + 2\pi_1 = 1$$

$$3\pi_1 = 1$$

$$\pi_1 = \frac{1}{3}$$

from (3), $\pi_2 = \frac{2}{3}$.

steady state dist - $\pi = \left[\frac{1}{3} \quad \frac{2}{3} \right]$.

2) If the initial state prob. distribution of a Markov chain is $p^{(0)} = \left(\frac{5}{6}, \frac{1}{6} \right)$ and the tpm of chain is $\begin{bmatrix} 0 & 1 \\ 1/2 & 1/2 \end{bmatrix}$ then find the prob. dist. of chain after 2 steps.

$$p^{(0)} = \left(\frac{5}{6}, \frac{1}{6} \right)$$

$$P = \begin{pmatrix} 0 & 1 \\ 1/2 & 1/2 \end{pmatrix}$$

after 2 steps

$$p^{(1)} = p^{(0)} \cdot P = \left(\frac{5}{6}, \frac{1}{6} \right) \begin{pmatrix} 0 & 1 \\ 1/2 & 1/2 \end{pmatrix}$$

$$p^{(1)} = \left(\frac{1}{12}, \frac{11}{12} \right)$$

$$P^{(2)} = P^{(1)} \cdot P = \begin{pmatrix} 1 & 1 \\ 12 & 12 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1/2 & 1/2 \end{pmatrix}$$

$$P^{(2)} = \begin{pmatrix} 11 & 13 \\ 24 & 14 \end{pmatrix}$$

Note:

The steady state distribution. is also called as limiting probabilities, probability in the long run invariant probabilities, stationary probabilities, fraction, proportion, How often.

2) A student's study habits are as follows. If he studies one night, he is 70% sure not to study next night. On the other hand the prob. that he does not study two nights in the succession is 0.6. In the long run, how often does he study.

S - study at night

T - not study at night.

$$P = \begin{matrix} & \begin{matrix} S & T \end{matrix} \\ \begin{matrix} S \\ T \end{matrix} & \begin{bmatrix} 0.3 & 0.7 \\ 0.4 & 0.6 \end{bmatrix} \end{matrix} \text{ Today.}$$

let $\pi = [\pi_1 \ \pi_2]$ be steady state dist,

$$\pi P = \pi \quad \text{where } \pi_1 + \pi_2 = 1 \rightarrow \textcircled{1}$$

$$[\pi_1 \ \pi_2] \begin{bmatrix} 0.3 & 0.7 \\ 0.4 & 0.6 \end{bmatrix} = [\pi_1 \ \pi_2]$$

$$[0.3\pi_1 + 0.4\pi_2 \quad 0.7\pi_1 + 0.6\pi_2] = [\pi_1 \ \pi_2]$$

compare same position elements,

$$0.3\pi_1 + 0.4\pi_2 = \pi_1$$

$$0.7\pi_1 + 0.6\pi_2 = \pi_2$$

$$-0.7\pi_1 + 0.4\pi_2 = 0,$$

$$0.7\pi_1 - 0.4\pi_2 = 0$$

$$-7\pi_1 + 4\pi_2 = 0$$

$$\Rightarrow \pi_1 - 4\pi_2 = 0 \rightarrow \textcircled{2}$$

$\rightarrow \textcircled{2}$

from (1), (2)

$$(2) \Rightarrow 7\pi_1 = 4\pi_2 \Rightarrow \pi_1 = \frac{4}{7}\pi_2$$

$$(1) \Rightarrow \pi_1 + \pi_2 = 1$$

$$\frac{4}{7}\pi_2 + \pi_2 = 1$$

$$4\pi_2 + 7\pi_2 = 7$$

$$11\pi_2 = 7$$

$$\pi_2 = \frac{7}{11}$$

$$\therefore \pi_1 = \frac{4}{7}\pi_2 = \frac{4}{7} \times \frac{7}{11} = \frac{4}{11}$$

$$\therefore \pi = \left[\frac{4}{11} \quad \frac{7}{11} \right]$$

3) The kpm is

3) Two boys B_1, B_2 and two girls G_1, G_2 are throwing a ball from one to another each boy throws the ball to other boy with prob. $\frac{1}{2}$ and to each girl with prob. $\frac{1}{4}$. On the other hand each girl throws the ball to each boy with prob. $\frac{1}{2}$ and never to other girl. In the long run how often does each receive the ball.

Boy \rightarrow other Boy prob. $\frac{1}{2}$
 \rightarrow each girl prob. $\frac{1}{4}$,

Girl \rightarrow each boy prob. $\frac{1}{2}$.
 \rightarrow never to other girl.

$$P = \begin{matrix} & \begin{matrix} B_1 & B_2 & G_1 & G_2 \end{matrix} \\ \begin{matrix} B_1 \\ B_2 \\ G_1 \\ G_2 \end{matrix} & \begin{bmatrix} 0 & \frac{1}{2} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{2} & 0 & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \end{bmatrix}_{4 \times 4} \end{matrix}$$

Let $\pi = [\pi_1 \quad \pi_2 \quad \pi_3 \quad \pi_4]$ be steady state dist. vector

$$\pi P = \pi$$

$$\text{E } \pi_1 + \pi_2 + \pi_3 + \pi_4 \rightarrow \textcircled{1}$$

$$\begin{bmatrix} \pi_1 & \pi_2 & \pi_3 & \pi_4 \end{bmatrix} \begin{bmatrix} 0 & 1/2 & 1/4 & 1/4 \\ 1/2 & 0 & 1/4 & 1/4 \\ 1/2 & 1/2 & 0 & 0 \\ 1/2 & 1/2 & 0 & 0 \end{bmatrix} = \begin{bmatrix} \pi_1 & \pi_2 & \pi_3 & \pi_4 \end{bmatrix}$$

$$\begin{bmatrix} \frac{\pi_2}{2} + \frac{\pi_3}{2} + \frac{\pi_4}{2} & \frac{\pi_1}{2} + \frac{\pi_2}{2} + \frac{\pi_4}{2} & \frac{\pi_1}{4} + \frac{\pi_2}{4} & \frac{\pi_1}{4} + \frac{\pi_2}{4} \end{bmatrix} = \begin{bmatrix} \pi_1 & \pi_2 & \pi_3 & \pi_4 \end{bmatrix}$$

compare same position element.

$$\frac{1}{2}(\pi_2 + \pi_3 + \pi_4) = \pi_1 \quad \frac{1}{2}(\pi_1 + \pi_2 + \pi_4) = \pi_2 \quad \frac{1}{4}(\pi_1 + \pi_2) = \pi_3 \quad \frac{1}{4}(\pi_1 + \pi_2) = \pi_4$$

$$-2\pi_1 + \pi_2 + \pi_3 + \pi_4 = 0 \quad \pi_1 - 2\pi_2 + \pi_3 + \pi_4 = 0 \quad \pi_1 + \pi_2 - 4\pi_3 = 0 \quad \pi_1 + \pi_2 - 4\pi_4 = 0$$

$$\pi_1 = 1/3, \pi_2 = 1/3, \pi_3 = 1/6, \pi_4 = 1/6.$$

Steady state distribution -

$$\pi = \begin{bmatrix} 1/3 & 1/3 & 1/6 & 1/6 \end{bmatrix}.$$

4) A Housewife buys 3 kinds of cereals A, B, C. she never buys the same cereal in successive weeks. If she buys cereal A, the next week she buys cereal B. However she buys B or C, next week she is 3 times as likely to buy A as other cereal. How often she buys each of the 3 cereals.

$$A \rightarrow B.$$

B \rightarrow 3 times as likely to buy A as other cereal C

$$C \rightarrow \text{'' ''}$$

$$P = \begin{matrix} & \begin{matrix} A & B & C \end{matrix} \\ \begin{matrix} A \\ B \\ C \end{matrix} & \begin{bmatrix} 0 & 1 & 0 \\ 3/4 & 0 & 1/4 \\ 3/4 & 1/4 & 0 \end{bmatrix} \end{matrix}$$

let $\pi = [\pi_1 \pi_2 \pi_3]$ be steady state distribution vector.

$$\pi P = \pi \quad \& \quad \pi_1 + \pi_2 + \pi_3 = 1 \rightarrow \textcircled{1}$$

$$[\pi_1 \quad \pi_2 \quad \pi_3] \begin{bmatrix} 0 & 1 & 0 \\ 3/4 & 0 & 1/4 \\ 3/4 & 1/4 & 0 \end{bmatrix} = [\pi_1 \quad \pi_2 \quad \pi_3]$$

$$\left[\frac{3\pi_2}{4} + \frac{3\pi_3}{4} \quad \pi_1 + \frac{\pi_2}{4} + \frac{\pi_3}{4} \quad \frac{\pi_2}{4} \right] = [\pi_1 \quad \pi_2 \quad \pi_3]$$

$$\frac{3\pi_2}{4} + \frac{3\pi_3}{4} = \pi_1, \quad \pi_1 + \frac{\pi_2}{4} = \pi_2, \quad \frac{\pi_2}{4} = \pi_3.$$

$$-4\pi_1 + 3\pi_2 + 3\pi_3 = 0 \rightarrow \textcircled{2} \quad 4\pi_1 - 4\pi_2 + \pi_3 = 0 \rightarrow \textcircled{3} \quad \pi_2 - 4\pi_3 = 0 \rightarrow \textcircled{4}$$

By solving $\textcircled{1}, \textcircled{2}, \textcircled{3}, \textcircled{4}$

$$P = 3/7 \quad \pi_1 = 3/7 \quad \pi_2 = \frac{16}{35} \quad \pi_3 = \frac{4}{35}$$

5) The transition prob. matrix of Markov chain $\{X_n\}, n=1, 2, 3, \dots$ having 3 states 1, 2 & 3 is

$$P = \begin{bmatrix} 0.1 & 0.5 & 0.4 \\ 0.6 & 0.2 & 0.2 \\ 0.3 & 0.4 & 0.3 \end{bmatrix} \text{ and the initial distribution } p^{(0)} = (0.7 \quad 0.2 \quad 0.1) \quad x_1=3$$

Find i) prob. $P(X_2=3)$ ii) $P(X_3=2, X_2=3, X_0=2)$

Given $p^{(0)} = (0.7 \quad 0.2 \quad 0.1)$

$$P(X_0=1) = 0.7 \quad P(X_0=2) = 0.2 \quad P(X_0=3) = 0.1$$

i) $P(X_2=3) = P(\text{at state 3})$

$$P^{(1)} = P^{(0)} \cdot P = [0.7 \quad 0.2 \quad 0.1] \begin{bmatrix} 0.1 & 0.5 & 0.4 \\ 0.6 & 0.2 & 0.2 \\ 0.3 & 0.4 & 0.3 \end{bmatrix}$$

$$P^{(1)} = \begin{bmatrix} 0.21 & 0.43 & 0.35 \end{bmatrix}$$

$$P^{(2)} = P^{(1)} \cdot P = [0.21 \quad 0.43 \quad 0.35] \begin{bmatrix} 0.1 & 0.5 & 0.4 \\ 0.6 & 0.2 & 0.2 \\ 0.3 & 0.4 & 0.3 \end{bmatrix}$$

$$= [0.385 \quad 0.336 \quad 0.279]$$

$$\therefore P(\text{at state 3}) = P(X_2=3) = 0.279$$

$$ii) P(X_3=2, X_2=3, X_1=3, X_0=2) = P(X_3=2|X_2=3) P(X_2=3|X_1=3) \\ P(X_1=3|X_0=2) \cdot P(X_0=2)$$

$$\Rightarrow P_{32}^{(1)} \cdot P_{33}^{(1)} \cdot P_{23}^{(1)} (0.2) \\ \Rightarrow (0.4)(0.3)(0.2)(0.2) \\ \Rightarrow 31625.$$

6) 3 Boys A, B, C are throwing a ball to each other. A always throw the ball to B and B always throw ball to C but C is just as likely to throw the ball to B as to A, show that the process is Markovian. Find the transition prob-matrix and classify the states.

A \rightarrow B

B \rightarrow C

C \rightarrow B
C \rightarrow A

$$P = \begin{matrix} & \begin{matrix} A & B & C \end{matrix} \\ \begin{matrix} A \\ B \\ C \end{matrix} & \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1/2 & 1/2 & 0 \end{bmatrix} \end{matrix}$$

future values depends on present values.

\therefore The chain is Markovian.

Irreducible :-

$$P^2 = P \cdot P = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1/2 & 1/2 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1/2 & 1/2 & 0 \end{bmatrix}$$

$$P^2 = \begin{bmatrix} 0 & 0 & 1 \\ 1/2 & 1/2 & 0 \\ 0 & 1/2 & 1/2 \end{bmatrix}$$

$$P^3 = P^2 \cdot P = \begin{bmatrix} 1/2 & 1/2 & 0 \\ 0 & 1/2 & 1/2 \\ 1/4 & 1/4 & 1/2 \end{bmatrix}$$

$$P^4 = P^3 \cdot P = \begin{bmatrix} 0 & 1/2 & 1/2 \\ 1/4 & 1/2 & 1/2 \\ 1/4 & 1/2 & 1/4 \end{bmatrix}$$

$$P^5 = P^4 \cdot P = \begin{bmatrix} 1/4 & 1/4 & 1/2 \\ 1/4 & 1/2 & 1/4 \\ 1/8 & 3/8 & 1/2 \end{bmatrix}$$

$$\therefore P_{11}^{(3)} > 0, P_{13}^{(2)} > 0, P_{21}^{(2)} > 0, P_{22}^{(2)} > 0, P_{33}^{(2)} > 0.$$

$$\therefore P_{ij}^{(n)} > 0 \text{ for some } n.$$

\therefore All states are irreducible.

\therefore Matrix is irreducible.

periodic:-

periodicity of state A:-

$$P_{11}^{(3)} > 0, P_{11}^{(5)} > 0, P_{11}^{(6)} > 0, \dots$$

$$\text{G.C.D} = \{3, 5, 6, \dots\} = 1 = d_i.$$

If $d_i = 1$ then state 'A' is a periodic.

periodic of B.

$$P_{22}^{(2)} > 0, P_{22}^{(3)} > 0, P_{22}^{(4)} > 0, \dots$$

$$\text{G.C.D} = \{2, 3, 4, \dots\} = 1$$

$$\therefore \text{period}(d_i) = 1$$

\therefore state B is aperiodic

\therefore all states are aperiodic.

state periodic of C

$$P_{33}^{(2)} > 0, P_{33}^{(3)} > 0, P_{33}^{(4)} > 0, \dots$$

$$\text{G.C.D} = \{2, 3, 4, \dots\} = 1$$

$$\therefore \text{period}(d_i) = 1$$

\therefore state C is aperiodic

\therefore All states are aperiodic


It has finite states.

The finite irreducible matrix becomes non-null, persistent.

All states are non-null, persistent & aperiodic.
 \therefore All states are ergodic.

(or)

Diagram:



The state A is reachable to state B & C

The state B " " " " C & A

" " C " " " A & B.

All states are reachable from all other states

\therefore The chain is irreducible.

It has infinite states.

\therefore finite irreducible matrix becomes non-null persistent.

period of A:-

G.C.D of $\{3, 5, 7, \dots\} = 1$.

\therefore state A is aperiodic

period of B:-

G.C.D $\{2, 3, 4, 5\} = 1$

\therefore state B is aperiodic

period of C:-

G.C.D $\{2, 3, 4, 5, \dots\} = 1$

state C is aperiodic

Note

1) Absorbing state:-


If a state i is called absorbing state if $P_{ii} = 1$.

2) If a matrix with absorbing state then it is not irreducible.

1) Construct the markov chain with transition prob matrix

$$P = \begin{bmatrix} A & 0.14 & 0.6 & 0 & 0 \\ B & 0.3 & 0.7 & 0 & 0 \\ C & 0.2 & 0.4 & 0.1 & 0.3 \\ D & 0 & 0 & 0 & 1 \end{bmatrix}$$


Is this matrix irreducible.



The state 4 is not reachable to state 1, 2, 3.
 \therefore The state 4 is absorbing state.

2) Find the nature of the states of the markov chain with the Gpm $\begin{bmatrix} 0 & 1 & 0 \\ 1/2 & 0 & 1/2 \\ 0 & 1 & 0 \end{bmatrix}$

diagraph:-



The state '0' is reachable from the states 1 & 2
 The state '1' is reachable from the states 0 & 2
 The state '2' is reachable from the states 1 & 0

every state is reachable from all other states,

\therefore The chain is irreducible

\therefore the chain is finite.

periodicity:-

period of state '0' \rightarrow G.C.D $\{2, 4, 6, 8, \dots\} = 2 = d_0$

period of '0' is $= d_0 = 2$.

period of state '1' :-

G.C.D $= \{2, 4, 6, \dots\} = 2 = d_1$

period of state '1' $= d_1 = 2$.

period of state '2' :-

G.C.D $= \{2, 4, 6, \dots\} = 2 = d_2$

\therefore period of state '2' is $= 2 = d_2$.

\therefore All states are non-null, persistent & aperiodic

\therefore All states are ergodic.

3) Suppose that the probability of dry day following rainy day is $1/2$ and the rainy day following is $1/2$ - Given that may 1st is dry day. Find that may 3rd is dry day and also may 5th is dry day.

Let D \rightarrow Dry day
R \rightarrow Rainy day.

$$P = \begin{matrix} & \begin{matrix} D & R \end{matrix} \\ \begin{matrix} D \\ R \end{matrix} & \begin{bmatrix} 1/2 & 1/2 \\ 1/3 & 2/3 \end{bmatrix} \end{matrix}$$

Given may 1st is dry day

$$P^{(0)} = \begin{bmatrix} D & R \\ 1 & 0 \end{bmatrix}$$

$$P^{(1)} = P^{(0)} \cdot P = \begin{bmatrix} 1 & 0 \\ 1/2 & 1/2 \\ 1/3 & 2/3 \end{bmatrix}$$

$$P^{(1)} = \begin{bmatrix} 1/2 & 1/2 \end{bmatrix}$$

$$P^{(2)} = P^{(1)} \cdot P = \begin{bmatrix} 1/2 & 1/2 \end{bmatrix} \begin{bmatrix} 1/2 & 1/2 \\ 1/3 & 2/3 \end{bmatrix}$$

$$P^{(2)} = \left[\frac{1}{4} + \frac{1}{6} \quad \frac{1}{4} + \frac{2}{6} \right] = \left[\frac{5}{12} \quad \frac{7}{12} \right]$$

$$P^{(3)} = P^{(2)} \cdot P = \left[\frac{5}{12} \quad \frac{7}{12} \right] \begin{bmatrix} 1/2 & 1/2 \\ 1/3 & 2/3 \end{bmatrix}$$

$$P^{(3)} = [0.402 \quad 0.597]$$

prob. of may 3rd day day = 0.402

$$P^{(4)} = P^{(3)} \cdot P$$

$$P^{(5)} = P^{(4)} \cdot P$$

41) A man either drives a car or catches the train to go to the office on each day, he never goes two days in a row by train but if he drives one day, the next day he is just as likely to drive again as he is to travel by train. Now suppose that, on the 1st day of week the man tossed a fair die and drove to work if 6 appear. Find:

i) prob. that he takes on the 3rd day.

ii) prob. that he drives to work in the long run.

Let C → car

T → Train

$$P = \begin{matrix} & \begin{matrix} T & C \end{matrix} \\ \begin{matrix} T \\ C \end{matrix} & \begin{bmatrix} 0 & 1 \\ 1/2 & 1/2 \end{bmatrix} \end{matrix}$$

He tossed a fair die & if it shows 5 or 6 then he goes by car.

$$P^{(1)} = \begin{matrix} & \begin{matrix} T & C \end{matrix} \\ \begin{matrix} T \\ C \end{matrix} & \begin{bmatrix} 5/6 & 1/6 \end{bmatrix} \end{matrix}$$

1) prob. of 3rd day by Train.

$$P^{(2)} = P^{(1)} \cdot P = \begin{bmatrix} \frac{5}{6} & \frac{1}{6} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix} = \begin{bmatrix} \frac{1}{12} & \frac{5}{6} + \frac{1}{12} \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{12} & \frac{11}{12} \\ 0 & 0 \end{bmatrix}$$

$$P^{(3)} = P^{(2)} \cdot P = \begin{bmatrix} \frac{1}{12} & \frac{11}{12} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix} = \begin{bmatrix} \frac{11}{24} & \frac{1}{12} + \frac{11}{24} \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} \frac{11}{24} & \frac{23}{24} \\ 0 & 0 \end{bmatrix}$$

Prob. of 3rd day goes by Train is $\frac{11}{24}$.